

Dissipative N – body code for galaxy evolution.

Peter Berczik & Sergei G. Kravchuk

Main Astronomical Observatory of
Ukrainian National Academy of Sciences
252650, Golosiiv, Kiev-022, Ukraine

e-mail: `berczik@mao.kiev.ua`

September 1, 1998

Abstract

The evolving galaxy is considered as a system of baryonic fragments embedded into the static dark nonbaryonic (DH) and baryonic (BH) halo and subjected to gravitational and viscous interactions. Although the chemical evolution of each separate fragment is treated in the frame of one – zone close box model with instantaneous recycling, its star formation (SF) activity is a function of mean local gas density and, therefore, is strongly influenced by other interacting fragments. In spite of its simplicity this model provides a realistic description of the process of galaxy formation and evolution over the Hubble timescale.

1 Introduction

Recent advances in extragalactic astrophysics show the close link of a disk galaxy dynamical evolution and its chemical and photometric behaviour over the Hubble timescale. In spite of remarkable succes of the modern theory of galaxy chemical evolution in explaining the properties of evolving galaxies ([4]) its serious shortcomings concern the multiparameter character and practical neglecting of dynamical effects. The inclusion of simplified dynamic into the chemical network ([6]) and vice versa the inclusion of simplified chemical scheme into the sophisticated 3D hydrodynamical code ([7, 1]) gives very promising results and allows to avoid a formal approach typical to standard theory.

In this paper the interplay between a disk galaxy dynamical evolution and its chemical behaviour is studied in a frame of a simplified model which provides a realistic description of the process of galaxy formation and evolution over the cosmological timescale.

2 Initial conditions

The evolving galaxy is treated as a system of baryonic fragments embedded into the extended halo composed of dark nonbaryonic and baryonic matter. The halo is modelled as a static structure with dark (DH) and diluted baryonic (BH) halo components having Plummer – type density profiles ([2]):

$$\rho_{BH}(r) = \frac{M_{BH}}{\frac{4}{3}\pi b_{BH}^3} \cdot \frac{b_{BH}^5}{(r^2 + b_{BH}^2)^{\frac{5}{2}}}$$

and

$$\rho_{DMH}(r) = \frac{M_{DMH}}{\frac{4}{3}\pi b_{DMH}^3} \cdot \frac{b_{DMH}^5}{(r^2 + b_{DMH}^2)^{\frac{5}{2}}},$$

where

$$M_{DMH} = 10^{12} M_{\odot}, \quad b_{DMH} = 25 \text{ kpc},$$

$$M_{BH} = 10^{11} M_{\odot}, \quad b_{BH} = 15 \text{ kpc}.$$

The dense baryonic matter (future galaxy disk and bulge) of total mass of $M_{gas} = 10^{11} M_{\odot}$ is assumed to be distributed among $N = 2109$ particles – fragments. The single particle density profile is also assumed to be a Plummer one. Its mass is taken to be $m_i = M_{gas}/N$ and radius $h_i = 1$ kpc. Initially all particles are smoothly placed inside the sphere of radius $R_{gas} = 50$ kpc and are involved into the Hubble flow ($H_0 = 75$ km/s/Mpc) and solid – body rotation around z axis. The initial motion of this system is described as:

$$\mathbf{V}(x, y, z) = [\boldsymbol{\Omega}(x, y, z) \times \mathbf{r}] + H_0 \cdot \mathbf{r} + \mathbf{DV}(x, y, z),$$

where $\boldsymbol{\Omega}(x, y, z) = (0, 0, 1) \cdot \Omega_{cir}$ is an angular velocity of the rotating sphere, $\Omega_{cir} = V_{cir}/R_{gas}$ and

$$V_{cir} = \sqrt{G \cdot \frac{M_{gas} + M_{DMH} + M_{BH}}{R_{gas}}}.$$

The components DV_x , DV_y , DV_z of the random velocity $\mathbf{DV}$ vector are assumed to be initially randomly distributed within an interval $0 \div 10$ km/s.

3 N – body code

The dynamical evolution of baryonic matter fragments which are subjected to gravitational influences of DM–baryonic halo and interfragment interactions is followed by means of effective N – body integrator with individual time step. The dynamics of such N – body system is described by following equations:

$$\begin{cases} d\mathbf{R}_i/dt = \mathbf{V}_i, \\ d\mathbf{V}_i/dt = \mathbf{A}_i(\mathbf{R}, \mathbf{V}). \end{cases} \quad (1)$$

The acceleration of i - th particle $\mathbf{A}_i$ is defined as a sum of three components.

$$\mathbf{A}_i = \mathbf{A}_i^{INT} + \mathbf{A}_i^{EXT} + \mathbf{A}_i^{VISC}, \quad (2)$$

where the first term $\mathbf{A}_i^{INT}$ accounts for gravitational interactions between fragments. The second one $\mathbf{A}_i^{EXT}$ is defined as an external gravitational acceleration caused by the DM and baryonic halo. The last term $\mathbf{A}_i^{VISC}$ corresponds to the viscous deceleration of fragment when passing through the baryonic halo.

The gravitational interaction between fragments is defined as the interaction of N Plummer profile elements:

$$\mathbf{A}_i^{INT} = -G \cdot \sum_{j=1, j \neq i}^N \frac{m_j}{(\mathbf{R}_{ij}^2 + h_{ij}^2)^{\frac{3}{2}}} \cdot \mathbf{R}_{ij}. \quad (3)$$

Here $h_{ij} = (h_i + h_j)/2$ and $\mathbf{R}_{ij} = \mathbf{R}_i - \mathbf{R}_j$.

Accounting for that the halo DM and baryonic components are also Plummer spheres the second term becomes:

$$\mathbf{A}_i^{EXT} = -G \cdot \left(\frac{M_{DMH}}{(\mathbf{R}_i^2 + b_{DMH}^2)^{\frac{3}{2}}} + \frac{M_{BH}}{(\mathbf{R}_i^2 + b_{BH}^2)^{\frac{3}{2}}} \right) \cdot \mathbf{R}_i. \quad (4)$$

The form of the last term $\mathbf{A}_i^{VISC}$ will be discussed in the next subsection.

The characteristic time step δt_i in the integration procedure for each particle is defined as:

$$\delta t_i = Const \cdot \min_j \left[\sqrt{\frac{|\mathbf{R}_{ij}|}{|\mathbf{A}_{ij}|}}, \frac{|\mathbf{R}_{ij}|}{|\mathbf{V}_{ij}|} \right] \quad (5)$$

where the $\mathbf{V}_{ij} = \mathbf{V}_i - \mathbf{V}_j$ and the $\mathbf{A}_{ij} = \mathbf{A}_i - \mathbf{A}_j$.

Here the $Const$ is a numerical parameter equal to $Const = 10^{-2}$ that provides a nice momentum and energy conservation over the integration interval of about 15 Gyr. For example, in the conservative case (when viscosity of the system is set equal to 0), the final total error in the energy equation is less than 1%.

4 Viscosity model

The viscosity term $\mathbf{A}_i^{VISC}$ is artificially introduced into the model so as to match the results of more sophisticated SPH approach on dynamical evolution of disk galaxies. The best fitness of results of this simplified approach with SPH modelling data (see *e.g.* [1]) is achieved when the momentum exchange between the baryonic halo and moving particles is modelled by the following expression:

$$\mathbf{A}_i^{VISC} = -k \cdot \mathbf{V}_i \cdot \frac{|\mathbf{V}_i|}{R_{VISC}} \cdot \frac{\rho_{BH}(r)}{\rho_{VISC}} \cdot \frac{m_i^{gas}}{m_i^{gas} + m_i^{star}}. \quad (6)$$

Here ρ_{VISC} and R_{VISC} are numerical parameters set equal to $\rho_{VISC} = 0.2 \text{ cm}^{-3}$ and $R_{VISC} = 10 \text{ kpc}$. The vector $\mathbf{V}_i$ is a particle velocity vector. It is to be noted that single particle is assumed to have a total mass which doesn't change with time and is defined as sum $m_i \equiv m_i^{gas} + m_i^{star}$. But masses of its gas and star components

m_i^{gas} and m_i^{star} are variable values and are defined by the temporal evolutionary status of the given fragment. Initially $m_i^{star} = 0$. The results of fitness show that for z component of viscosity term $k = 1$. In the galaxy plane where it is necessary to account for baryonic halo and baryonic fragments partial corrotation the dynamical friction is decreased and for x, y component of viscosity term k reduced to the value 0.15.

5 Density definition

In the frame of the multifragmented model the definition of local gas density is introduced in the SPH manner, e.g. local gas density depends on the total mass of matter contained in the sphere of radius H_i around the i -th particle. For each i -th particle the value of its smoothing radius H_i is choosed (using the quicksort algorithm) requiring that the volume of such radius comprises $N_B = 21$ nearest particles (i.e. $\approx 1\%$ of total number of particles N). Therefore, the total mass M_i and density of gas ρ_i inside this sphere are defined as

$$M_i = \sum_{j=1}^N \Delta m_{ij}^{gas}, \quad \rho_i = \frac{M_i}{\frac{4}{3}\pi H_i^3}, \quad (7)$$

where Δm_{ij}^{gas} is defined as

$$\begin{aligned} \text{if } |\mathbf{R}_{ij}| > (H_i + h_j) &\Rightarrow \Delta m_{ij}^{gas} = 0, \\ \text{if } |\mathbf{R}_{ij}| < (H_i - h_j) &\Rightarrow \Delta m_{ij}^{gas} = m_j^{gas}, \\ \text{else } \Delta m_{ij}^{gas} &= m_j^{gas} \cdot \frac{H_i + h_j - |\mathbf{R}_{ij}|}{2 \cdot h_j}. \end{aligned}$$

6 Star formation and SN explosions

A forming disk galaxy is modelled as a system of interacting fragments (named as particles) embedded into the extended halo. Each particle is composed of gas and stellar components and its total mass is defined as $m_i \equiv m_i^{gas} + m_i^{star}$. Initially all particles are purely gaseous and, therefore, initially $m_i^{star} = 0$. To follow a particle star formation (SF) activity a special timemark t_i^{begSF} is introduced which initially is set equal to $t_i^{begSF} = 0$. The particles eligible to star formation events are chosen as particles which still have a sufficient amount of the gas component and their densities exceed some critical value ρ_{minSF} during some fixed time interval Δt_{SF} (of order of free-fall time):

$$\left\{ \begin{array}{l} \rho_i > \rho_{minSF}, \\ t_i - t_i^{begSF} > \Delta t_{SF}, \\ m_i^{gas} / (m_i^{gas} + m_i^{star}) > 10^{-4}. \end{array} \right. \quad (8)$$

Here $\Delta t_{SF} = 50$ Myr, and $\rho_{minSF} = 0.01 \text{ cm}^{-3}$ (this last value is not crucial and is only limiting one).

If the particle was subjected to SF activity the parameter t_i^{begSF} is set equal to $t_i^{begSF} = t_i$, and m_i^{star} and m_i^{gas} are redefined as

$$\begin{cases} m_i^{star} := \epsilon \cdot (1 - R) \cdot m_i^{gas} + m_i^{star}, \\ m_i^{gas} := (1 - \epsilon \cdot (1 - R)) \cdot m_i^{gas}. \end{cases} \quad (9)$$

Here ϵ is a SF efficiency which is defined as

$$\epsilon = \alpha \cdot \frac{\rho_i}{\rho_{SF}} \cdot (1 - \exp(-\frac{\rho_{SF}}{\rho_i})). \quad (10)$$

Therefore,

$$\begin{aligned} \text{if } \frac{\rho_i}{\rho_{SF}} \rightarrow 0 &\Rightarrow \epsilon \rightarrow \alpha \cdot \frac{\rho_i}{\rho_{SF}}, \\ \text{if } \frac{\rho_{SF}}{\rho_i} \rightarrow 0 &\Rightarrow \epsilon \rightarrow \alpha. \end{aligned}$$

To match available observational data on star formation efficiency (see *e.g.* [8]) parameters α and ρ_{SF} are set equal to $\alpha = 0.5$ and $\rho_{SF} = 10 \text{ cm}^{-3}$. The chemical evolution of each separate fragment is treated in the frame of one - zone close box model with instantaneous recycling. In the frame of this approach for the returned fraction of gas from evolved stars is taken a standard value $R = 0.25$.

Following the instantenuous recycling approximation it is assumed that after each SF and SN explosions the heavy element enriched gas is returned to the system and mixed with old (heavy element deficient) gas. After each act of SF and SN explosions the value of heavy element abundances of gas in particle is upgraded according to:

$$Z_i := Z_i + \frac{\epsilon \cdot R \cdot \Delta Z}{1 - \epsilon \cdot (1 - R)}. \quad (11)$$

The value $\Delta Z = 0.01$ is used as an average value for all values of $Z = 0.001 \div 0.04$ (see [1]). Initially $Z_i = 0.0$ in all particles.

7 Conclusion

The proposed simple model provides the self - consistent picture of the process of galaxy formation, its dynamical and chemical evolution is in agreement with the results of more sophisticated approaches (see *e.g.* [7, 5, 6, 1]).

- The rapidly rotating protogalaxy finally formed a three - component system resembling a typical spiral galaxies: a thin disk and spheroidal component made of gas and/or stars and a dark matter halo.

Fig.1 and Fig.2 show respectively the star formation rate and the galaxy total stellar mass as a function of time. Fig.3 shows the cylindrical distribution of stellar (upper curve) and gaseous (lower one) components of the final model disk galaxy as a function of a distance from the galaxy center in the galactic plane.

The total star formation rate (SFR) is a succession of short bursts which doesn't exceed $28 M_{\odot}/\text{yr}$. During first 2 Gyr of evolution only about 20% of total galaxy mass is transformed into the stars. The SFR gradually decreases, during the further evolution, to the value of about $2 M_{\odot}/\text{yr}$ typical for our own Galaxy. The final total stellar and gas mass of the model galaxy disk are about 92% and 8%. All these data as well as surface densities distributions of stellar and gaseous components (see Fig.4) are in nice agreement with present date observational data ([3, 4]).

- The disk component possesses a typical spiral galaxy rotation curve and the distribution of radial and V_z - th velocities of baryonic particles clearly show the presence of the central bulge (see Fig.6 and Fig.7).
- The metallicities and the global metallicity gradient resemble distributions observed in our own Galaxy (Fig.5). The averaged observed value of global metallicity $Z/Z_{\odot}(r)$ (see [4]) is shown in this Fig. as a solid line.

Acknowledgements: Peter Berczik would like to acknowledge the American Astronomical Society for financial support of this work under International Small Research Grant.

8 Figures

Figure 1: The variation of total star formation rate of forming disk galaxy with time.

Figure 2: The growth of galaxy stellar mass with time.

Figure 3: The cylindrycal distribution of masses of stellar (upper curve) and gaseous (lower one) components as a function of distance from galaxy center in a galactic plane.

Figure 4: The surface density radial distributions of stellar and gaseous components (stellar component is shown by filled stars, gaseous one by astericks). Theoretical distributions (not scaled) of surface density for radial exponential scale lengths 2.0 and 3.0 kpc are shown below by lines.

Figure 5: The radial distribution of heavy element abundances (averaged observed distribution of z for our Galaxy is shown by solid line).

Figure 6: The final distribution of V_z - th velocities of baryonic gas–stellar particles.

Figure 7: The final galaxy rotation curve.

References

- [1] Berczik P., Kravchuk S.G., 1997, Ap.Sp.Sci. **245**, 27
- [2] Douphole B., Colin J., 1995, A.&A. **300**, 117
- [3] Kuijken K. Gilmore G., 1989, M.N.R.A.S. **239**, 605
- [4] Pagel B.E.J., 1994, in "The Formation and Evolution of Galaxies", Eds. Munoz - Tunon, C., Sanchez, F., Cambridge Univ. Press
- [5] Raiteri C.M., Villata M., Navarro J.F., 1996, A.&A. **315**, 105
- [6] Samland M., Hensler G., 1996, Reviews in Modern Astronomy **9**, 277
- [7] Steinmetz M., Muller E., 1994, A.&A. **281**, L97
- [8] Wilking B.A., Lada C.J., 1983, Ap.J. **274**, 698









